

Response of Birds to Fire in the American Southwest¹

Carl E. Bock² and William M. Block³

Abstract

Fire was a common prehistoric disturbance in most southwestern grasslands, oak savannas, and coniferous forests, but not in Sonoran and Mojave desertscrub, or in riparian ecosystems. Prescribed burning should be applied, but under experimental conditions that facilitate studying its impacts on birds and other components of biodiversity. Fire plays a critical role in maintaining a balance between desert grassland and Chihuahuan desertscrub, but unburned areas also are important for birds dependent upon woody vegetation and/or heavy grass cover. Understory fire probably once played a critical role in maintaining relatively open oak (*Quercus* spp.), pinyon-juniper (*Pinus-Juniperus*), and ponderosa pine (*Pinus ponderosa*) woodlands and their bird assemblages, while stand replacement fires sustained aspen groves (*Populus tremuloides*) at higher elevations. Carefully controlled prescribed burning, thinning, and grazing management will be needed to return fire to its prehistoric role in these habitats. There is an urgent need for cooperative effort between managers and researchers to implement replicated burns to quantify avian responses in appropriate habitats.

Key words: birds, desert, fire, grassland, mixed-conifer, pine-oak, prescribed burning, riparian, savanna, Southwest.

The ecological importance of fire in the American Southwest has long been recognized (Leopold 1924), but the nature of these fires has been altered drastically by disturbances such as logging, livestock grazing, and suppression efforts (Moir et al. 1997, McPherson and Weltzin 2000). Southwestern ecosystems range from low desert to high montane, and fire ecology and man-

agement issues differ greatly among them (Krammes 1990).

The Sonoran and Mojave Deserts consist of structurally complex vegetation (Turner et al. 1995), and they have correspondingly rich avifaunas dependent upon trees and large succulents (Mills et al. 1991). The spread of exotic grasses such as red brome (*Bromus rubens*) and buffelgrass (*Pennisetum ciliare*) have increased the frequency and intensity of fire in these deserts, killing much of the native vegetation (Miller et al. 1995). Fire effects on birds in these deserts therefore are almost entirely negative.

Most southwestern grasslands were invaded by woody plants from the Chihuahuan Desert, beginning especially in the late 1800s, largely because of decreased frequencies and intensities of fire (Archer et al. 1995, McPherson 1995). Birds associated with grasslands have declined more than other avian groups, both nationally and in the Southwest (Brown and Davis 1998, Vickery and Herkert 2001), at least in part because of the conversions of former grassland to desertscrub. Fire has the short term effect in southwestern grasslands of reducing grass cover while stimulating forb cover and seed production for two or three post-fire growing seasons. Such burning favors grassland birds associated with relatively open ground, such as Mourning Dove (*Zenaida macroura*), Horned Lark (*Eremophila alpestris*), and Lark Sparrow (*Chondestes grammacus*), over those requiring heavy grass cover, such as Grasshopper Sparrow (*Ammodramus savannarum*), Botteri's Sparrow (*Aimophila botterii*), and Cassin's Sparrow (*Aimophila cassinii*; fig. 1). Prescription fire should be applied in these ecosystems, especially in relatively mesic sites, to restore open grasslands and retard woody encroachment (Bock and Bock 1992, Gordon 2000). However, a balanced approach is called-for, since other species prefer or require woody cover in Chihuahuan shrub/grasslands (Whitford 1997, Pidgeon et al. 2001).

Interior chaparral occurs from northwestern Arizona to southwest Texas and northern Mexico, and supports a variety of shrubland birds (Szaro 1981, Pase and Brown 1982). Drought, livestock grazing, and fire suppression have increased shrub cover and reduced cover of perennial grasses and forbs in Arizona chaparral (Brejda 1997). Virtually nothing is known about avian

¹A version of this paper was presented at the **Third International Partners in Flight Conference, March 20-24, 2002, Asilomar Conference Grounds, California.**

²Department of Environmental, Population, and Organismic Biology, University of Colorado, Boulder, CO 80309-0334 email: carl.bock@colorado.edu

³USDA Forest Service, Rocky Mountain Research Station, 2500 S. Pine Knoll, Flagstaff, AZ 86001.

responses to fire in this ecosystem (but see Szaro 1981), and experiments with prescribed fire are strongly recommended.




Figure 1— Relative abundances of four grassland birds two years before, just following, and for three summers after fire in a semi-arid grassland in southeastern Arizona (data from Bock and Bock 1992).

Fire once maintained oak (*Quercus* spp.) savannas in the Mexican borderlands in an open condition, by frequently killing smaller trees and shrubs (McPherson 1997). These ecosystems became more densely wooded when livestock grazing and fire suppression reduced fine fuels (McPherson and Weltzin 2000). Typical birds of southwestern oak savannas include species specializing on acorns, insect hawkers, foliage gleaners, and species dependent on the grassy understory (Brown 1982). Responses of these birds to prescribed or wildfire in southwestern oak woodlands are largely unknown, but studies in Midwestern savannas suggest that fire would favor insect hawkers and understory species over foliage gleaners (Davis et al. 2000).

Southwestern riparian woodlands support an abundance and variety of breeding birds vastly greater than adjacent ecosystems (Carothers et al. 1974), but many of these woodlands have been damaged or completely eliminated by anthropogenic forces such as alteration of natural flooding regimes and the spread of exotic

vegetation (Ellis et al. 1998). The result has been a substantial decline in abundance and variety of birds in these ecosystems (Rosenburg et al. 1991), although remnant patches continue to support high avian biodiversity (Strong and Bock 1990, Skagen et al. 1998). There is little evidence that fire was a common historical event in southwestern riparian woodlands, perhaps because of moisture. Recent fires have been highly destructive of native riparian vegetation, most particularly the large trees such as cottonwood (*Populus* spp.) and sycamore (*Platanus* spp.) that are especially important to birds (Bock and Bock 1984, Busch 1995). Therefore, prescribed fire is not recommended for these ecosystems.

Pinyon-juniper (*Pinus-Juniperus* spp) woodland is widespread throughout the Southwest, but little is known about fire effects in this ecosystem (Balda and Masters 1980, Severson and Rinne 1990). We found no studies related to birds and fire in pinyon juniper woodlands. Fire likely would favor species associated with more open habitats, while negatively impacting species that depend on trees for foraging or nesting, but this tenuous conclusion is based on responses of birds to mechanical clearing of dense stands rather than to fire itself (Sedgwick and Ryder 1987).

Fire is perhaps the most important natural disturbance in southwestern ponderosa pine (*Pinus ponderosa*) forests, and frequent low-intensity fires were part of their evolutionary history (Pyne 1996, Moir et al. 1997). Suppression efforts, livestock grazing, and logging have altered natural fire regimes, resulting in dense forests and increased risk of unnatural stand replacement fires. Most studies of fire effects on birds in these forests have focused on stand replacement burns, where woodpeckers and ground foragers are the main short-term beneficiaries (Dwyer and Block 2000). Species likely to benefit from cooler understory fires are those requiring relatively open habitats, such as flycatchers, swallows, and bluebirds (Marshall 1963, Horton and Mannan 1988).

Bird numbers and species richness seemed to respond positively shortly after the 1996 Horseshoe and Hochderffer fires on the Coconino National Forest Arizona (table 1). During the breeding season three years post-fire, more species were detected in areas where fires were severe (stand-replacement) and moderate (understory) than in adjacent unburned forests (45, 41, and 31 species, respectively). A similar trend was found during the nonbreeding season with 33, 35, and 26 species detected in severe, moderate, and unburned forests. Major species groups that increased in response to fire included woodpeckers, flycatchers, and thrushes. In contrast, many foliage-gleaning birds (Mountain Chickadees, Plumbeous Vireo, Pygmy Nuthatch, Yellow-rumped Warbler, and Grace's Warbler) were detected less frequently within severe fire areas.

Table 1— Simple counts of birds detected during point counts conducted within ponderosa pine forests burned during the 1996 Horseshoe and Hochdjerfer fires, Coconino National Forest, Arizona (Block, unpubl. data). Counts were conducted in the breeding season of 1999 and nonbreeding season 1999–2000, approximately three years after the fires.

Species	Breeding			Nonbreeding		
	Severe	Moderate	Unburned	Severe	Moderate	Unburned
Golden Eagle				1		
Bald Eagle				1	1	1
Sharp-shinned hawk	1			7	1	
Cooper's Hawk		1				
Northern Goshawk			1			2
Red-tailed Hawk	1	3		3	1	
American Kestrel	1	3		3		
Merlin					1	
Wild Turkey				2		
Mourning Dove	37			1	5	
Long-eared Owl		32	30		1	
Northern Pygmy Owl				6		
Common Nighthawk		2				
Broad-tailed Hummingbird	20	34	5			
Rufous Hummingbird		1				
Northern Flicker	32	28	17	103	83	42
Acorn Woodpecker	6					
Lewis's Woodpecker	9					
Williamson's Sapsucker				1	12	4
Yellow-bellied Sapsucker					1	
Hairy Woodpecker	88	28	10	149	46	29
Three-toed Woodpecker	1			11		
Red-naped Sapsucker						
Cordilleran Flycatcher	1		1		1	3
Olive-sided Flycatcher	12	1				
Western Wood-pewee	51	93	11			
Horned Lark	1					
Violet-green Swallow		3	10			
Pinyon Jay	4	36	17	22	23	2
Steller's Jay	33	94	88	152	189	114

Table 1—contd.

Species	Breeding			Nonbreeding		
	Severe	Moderate	Unburned	Severe	Moderate	Unburned
Clark's Nutcracker	3	4		39	44	22
American Crow	5		3	15	16	20
Common Raven	11	5	8	96	87	83
Mountain Chickadee	2	74	111	11	117	104
Bushtit				4	7	6
Brown Creeper	8	10	6	3	25	23
White-breasted Nuthatch	43	65	76	63	121	95
Red-breasted Nuthatch				2	1	
Pygmy Nuthatch	1	26	26	17	130	116
House Wren	2	1				
Rock Wren		1				
Golden-crowned Kinglet					3	1
Ruby-crowned Kinglet	144	136	77	116	106	74
Western Bluebird	6			4	3	1
Mountain Bluebird	1	5	17	1	6	4
Townsend's Solitaire	3					
Hermit Thrush	53	46	17	35	43	13
American Robin	10	36	24		1	
Plumbeous Vireo		1				
Warbling Vireo			3			
Virginia's Warbler	10	69	78	1		
Yellow-rumped Warbler		31	32			1
Grace's Warbler			2			2
Olive Warbler	3	5				
Black-headed Grosbeak	2		4			
Vesper Sparrow	6					
Lark Sparrow	19	51	9	1	4	
Chipping Sparrow	31	89	60	71	94	46
Dark-eyed Junco	35	14		1	2	
Western Meadowlark	1	13	2			
Brown-headed Cowbird	24	103	20			
Western Tanager						
Summer Tanager		3				

Table 1—contd.

Species	Breeding			Nonbreeding		
	Severe	Moderate	Unburned	Severe	Moderate	Unburned
Hepatic Tanager	1					
Pine Siskin	3	3	3	11	2	5
Lesser Goldfinch	1	1				
Red Crossbill	4	4	7	14	30	54
House Finch	2	1				
Evening Grosbeak	2		5		1	1
Total number of species	45	41	31	33	35	26
Total number of detections	734	1160	779	967	1209	867

^aN = 49 counting stations within severe, moderate, and unburned areas. Counting stations were sampled three times during the breeding season and four times during the nonbreeding season.

Mixed conifer forests of the Southwest are dominated by Douglas-fir (*Pseudotsuga menziesii*), white fir (*Abies concolor*), Engelmann spruce (*Picea engelmannii*), and aspen (*Populus tremuloides*; Brown 1982). Many fires that occur at higher elevations in these forests are stand replacing, providing opportunities for establishment of aspen (Moir 1993), and the variety of birds associated with it (Finch and Reynolds 1987, Hutto 1995). However, a variety of other birds, including the threatened Mexican Spotted Owl (*Strix occidentalis lucida*), depend on substantial conifer forests (Smith 1980, U.S. Department of Interior 1995). More field research specific to Southwestern conditions needs to be conducted to understand fire effects on birds in mixed-conifer forests.

Both deliberately-set prescribed fire and prescribed natural fire (natural ignition, allowed to burn within pre-set limits) are essential approaches in reducing fuels and returning southwestern grass/shrublands, oak savannas, and conifer forests to their prehistoric conditions. Given the historical importance of fire in these ecosystems, there have been surprisingly few studies of the effects of either wildfire or prescribed burning on southwestern bird populations and communities. We call for managers and researchers to cooperate in the design and implementation of replicated, large-scale, properly controlled field experiments to examine avian responses to prescription burning.

Acknowledgments

We thank Victoria Saab for organizing the session on fire and birds, of which this paper was a part.

Literature Cited

Archer, S., D. S. Scheme, and E. A. Holland. 1995. **Mechanisms of shrubland expansion: land use, climate or CO₂?** Climatic Change 29: 91-99.

Balda, R. P., and N. Masters. 1980. **Avian communities in the pinyon-juniper woodland.** In: R. M. De-Graff, technical coordinator. Workshop proceedings, management of western forests and grasslands for nongame birds. Gen. Tech. Rep. INT-86. Ogden, UT: Intermountain Research Station, Forest Service, U.S. Department of Agriculture; 146-167.

Bock, C. E., and J. H. Bock. 1984. **Importance of sycamores to riparian birds in southeastern Arizona.** Journal of Field Ornithology 55: 97-103.

Bock, C. E., and J. H. Bock. 1992. **Response of birds to wildfire in native versus exotic Arizona grassland.** Southwestern Naturalist 37: 73-81.

- Brejda, J. J. 1997. **Soil changes following 18 years of protection from grazing in Arizona chaparral.** Southwestern Naturalist 42: 478-487.
- Brown, D. E. (editor). 1982. **Biotic communities of the American Southwest - United States and Mexico.** Desert Plants 4: 1-342.
- Brown, D. E., and R. Davis. 1998. **Terrestrial bird and mammal distribution changes in the American Southwest, 1890-1990.** In: B. Tellman, D. M. Finch, C. Edminster, and R. Hamre, editors. The future of arid grasslands: identifying issues, seeking solutions. Forest Service Proceedings RMRS-P-3. Fort Collins, CO: Rocky Mountain Research Station, Forest Service, U.S. Department of Agriculture; 47-64.
- Busch, D. E. 1995. **Effects of fire on southwestern riparian plant community structure.** The Southwestern Naturalist 40: 259-267.
- Carothers, S. W., R. R. Johnson, and S. W. Aitchison. 1974. **Population structure and social organization of southwestern riparian birds.** American Zoologist 14: 97-108.
- Davis, M. A., D. W. Peterson, P. B. Reich, M. Crozier, and T. Query. 2000. **Restoring savanna using fire: impact on the breeding bird community.** Restoration Ecology 8: 30-40.
- Dwyer, J. K., and W. M. Block. 2000. **Effects of wildfire on densities of secondary cavity nesting birds in ponderosa pine forests of northern Arizona.** Tall Timbers Fire Ecology Conference Proceedings 21: 151-156.
- Ellis, L. M., C. S. Crawford, and M. C. Molles. 1998. **Comparison of litter dynamics in native and exotic riparian vegetation along the Middle Rio Grande of central New Mexico, USA.** Journal of Arid Environments 38: 283-296.
- Finch, D. M., and R. T. Reynolds. 1987. **Bird response to understory variation and conifer succession in aspen forests.** In: J. Emerick, S. Q. Foster, L. Hayden-Wing, J. Hodgson, J. W. Monarch, A. Smith, O. Thorne II, and J. Todd, editors. Issues and technology in the management of impacted wildlife. Boulder, CO: Thorne Ecological Institute; 87-96.
- Gordon, C. E. 2000. **Fire and cattle grazing on wintering sparrows in Arizona grasslands.** Journal of Range Management 53: 384-389.
- Horton, S. P., and R. W. Mannan. 1988. **Effects of prescribed fire on snags and cavity-nesting birds in southeastern Arizona pine forests.** Wildlife Society Bulletin 16: 37-44.
- Hutto, R. L. 1995. **Composition of bird communities following stand-replacement fires in northern Rocky Mountains.** Conservation Biology 9: 1041-1058.
- Krammes, J. S., technical coordinator. 1990. **Effects of fire management of southwestern natural resources.** Gen. Tech. Rep. RM-191. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture; 293 p.
- Leopold, A. 1924. **Grass, brush, and timber fire in southern Arizona.** Journal of Forestry 22: 1-10.
- Marshall, J. T., Jr. 1963. **Fire and birds in the mountains of southern Arizona.** Proceedings of the Tall Timbers Fire Ecology Conference 2: 134-141.
- McPherson, G. R. 1995. **The role of fire in desert grasslands.** In: M. P. McClaran and T. R. Van Devender, editors. The desert grassland. Tucson, AZ: University of Arizona Press; 130-151.
- McPherson, G. R. 1997. **Ecology and management of North American savannas.** Tucson, AZ: University of Arizona Press; 208 p.
- McPherson, G. R., and J. F. Weltzin. 2000. **Disturbance and climate change in United States/Mexico borderland communities.** Gen. Tech. Rep. RMRS-GTR-50. Fort Collins, CO: Rocky Mountain Research Station, Forest Service, U. S. Department of Agriculture; 24 p. Miller, M., C. Gossard, and K. Mahoney. 1995. Fire management consideration for BLM Wilderness Areas. In: J. K. Brown, R. W. Mutch, C. W. Spoon, and R. H. Wakimoto, technical coordinators. Proceedings: symposium on fire in wilderness and park management. Gen. Tech. Rep. INT-GTR-320. Ogden, UT: Intermountain Research Station, Forest Service, U.S. Department of Agriculture; 230-232.
- Mills, G. S., J. B. Dunning Jr., and J. M. Bates. 1991. **The relationship between breeding bird density and vegetation volume.** Wilson Bulletin 103: 468-479.
- Moir, W. H. 1993. **Alpine tundra and coniferous forest.** In: W. A. Dick-Peddie, editor. New Mexico vegetation: past, present, and future. Albuquerque, NM: University of New Mexico Press; 47-84.
- Moir, W. H., B. W. Geils, M. A. Benoit, and D. Scurlock. 1997. **Ecology of Southwestern ponderosa pine forests.** In: W. M. Block and D. M. Finch, technical editors. Songbird ecology in Southwestern ponderosa pine forests: a literature review. Gen. Tech. Rep. RM-GTR-292. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture; 3-27.
- Pase, C. P., and D. E. Brown. 1982. **Interior chaparral.** Desert Plants 4: 95-99.
- Pidgeon, A. M., N. E. Mathews, R. Benoit, and E. V. Nordheim. 2001. **Response of avian communities to historic habitat change in the northern Chihuahuan Desert.** Conservation Biology 15: 1772-1788.
- Pyne, S. J. 1996. **Nouvelle Southwest.** In: W. W. Covington and P. K. Wagner, technical coordinators. Conference on adaptive ecosystem restoration and management: restoration of cordilleran conifer landscapes of North America. Gen. Tech. Rep. RM-GTR-278. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture; 10-16.
- Rosenberg, K. V., R. D., Ohmart, W. C. Hunter, and B. W. Anderson. 1991. **Birds of the lower Colorado River valley.** Tucson, AZ: University of Arizona Press; 416 p.
- Sedgwick, J. A., and R. A. Ryder. 1987. **Effects of chaining on nongame wildlife.** In: R. L. Everitt, compiler. Proceedings

Fire and Southwestern Birds – Bock and Block

- of the pinyon-juniper conference. Gen. Tech. Rep. INT-215. Ogden, UT: Intermountain Research Station, Forest Service, U.S. Department of Agriculture; 541-551.
- Severson, K. E., and J. N. Rinne. 1990. **Increasing habitat diversity in Southwestern forests and wood-lands via prescribed burning.** In: J. S. Krammes, technical coordinator. Effects of fire management of south-western natural resources. Gen. Tech. Rep. RM-191. Fort Collins, CO: Rocky Mountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture; 94-104.
- Skagen, S. K., C. P. Melcher, W. H. Howe, and F. L. Knopf. 1998. **Comparative use of riparian corridors and oases by migrating birds in southeast Arizona.** Conservation Biology 12: 896-909.
- Smith, K. G. 1980. **Nongame birds of the Rocky Mountain spruce-fire forests and their management.** In: R. M. DeGraff, technical coordinator. Workshop proceedings, management of western forests and grasslands for nongame birds. Gen. Tech. Rep. INT-86. Ogden, UT: Intermountain Research Station, Forest Service, U.S. Department of Agriculture; 258-279.
- Strong, T. R., and C. E. Bock. 1990. **Bird species distribution patterns in riparian habitats in southeastern Arizona.** Condor 92: 866-885.
- Szaro, R. C. 1981. **Bird population responses to converting chaparral to grassland and riparian habitats.** Southwestern Naturalist 26: 251-256.
- Turner, R. M., J. E. Bowers, and T. L. Burgess. 1995. **Sonoran Desert plants.** Tucson, AZ: University of Arizona Press; 504 p.
- U. S. Department of Interior. 1995. **Recovery plan for the Mexican Spotted Owl (*Strix occidentalis lucida*).** Volume 1. Albuquerque, NM: Fish and Wildlife Service, U.S. Department of Interior.
- Vickery, P. D., and J. R. Herkert. 2001. **Recent advances in grassland bird research: where do we go from here?** Auk 118: 11-15.
- Whitford, W. G. 1997. **Desertification and animal biodiversity in the desert grasslands of North America.** Journal of Arid Environments 37: 709-720.